Package 'breastCancerUNT'

June 5, 2025

Type Package Title Gene expression dataset published by Sotiriou et al. [2007] (UNT). **Version** 1.46.0 Date 2011-02-10 Description Gene expression data from a breast cancer study published by Sotiriou et al. in 2007, provided as an eSet. biocViews ExperimentData, CancerData, BreastCancerData, MicroarrayData, TwoChannelData, GEO Author Markus Schroeder, Benjamin Haibe-Kains, Aedin Culhane, Christos Sotiriou, Gianluca Bontempi, John Quackenbush Maintainer Markus Schroeder <mschroed@jimmy.harvard.edu>, Benjamin Haibe-Kains <bhaibeka@jimmy.harvard.edu> **Depends** R (>= 2.5.0) Suggests survcomp, genefu, Biobase LazyLoad yes License Artistic-2.0 URL http://compbio.dfci.harvard.edu/ git_url https://git.bioconductor.org/packages/breastCancerUNT git_branch RELEASE_3_21 git_last_commit 126eb96 git_last_commit_date 2025-04-15 **Repository** Bioconductor 3.21 Date/Publication 2025-06-05

Contents

4

Index

Gene expression, annotations and clinical data from Sotiriou et al. 2006

Description

This dataset contains the gene expression, annotations and clinical data as published in Sotiriou et al. 2006.

Usage

data(unt)

Format

ExpressionSet with 44928 features and 137 samples, containing:

- exprs(unt): Matrix containing gene expressions as measured by Affymetrix hgu133a/hgu133b technology (single-channel, oligonucleotides).
- fData(unt): AnnotatedDataFrame containing annotations of Affy microarray platform hgu133a and hgu133b.
- pData(unt): AnnotatedDataFrame containing Clinical information of the breast cancer patients whose tumors were hybridized.
- experimentalData(unt): MIAME object containing information about the dataset.
- annotation(unt): Name of the affy chip.

Details

This dataset represent the study published by Sotiriou et al. 2006.

• Abstract: Background: Histologic grade in breast cancer provides clinically important prognostic information. However, 30%-60% of tumors are classified as histologic grade 2. This grade is associated with an intermediate risk of recurrence and is thus not informative for clinical decision making. We examined whether histologic grade was associated with gene expression profiles of breast cancers and whether such profiles could be used to improve histologic grading. Methods: We analyzed microarray data from 189 invasive breast carcinomas and from three published gene expression datasets from breast carcinomas. We identified differentially expressed genes in a training set of 64 estrogen receptor (ER)-positive tumor samples by comparing expression profiles between histologic grade 3 tumors and histologic grade 1 tumors and used the expression of these genes to define the gene expression grade index. Data from 597 independent tumors were used to evaluate the association between relapse-free survival and the gene expression grade index in a Kaplan-Meier analysis. All statistical tests were two-sided. Results: We identified 97 genes in our training set that were associated with histologic grade; most of these genes were involved in cell cycle regulation and proliferation. In validation datasets, the gene expression grade index was strongly associated with histologic grade 1 and 3 status; however, among histologic grade 2 tumors, the index spanned the values for histologic grade 1-3 tumors. Among patients with histologic grade 2 tumors, a high

unt

gene expression grade index was associated with a higher risk of recurrence than a low gene expression grade index (hazard ratio = 3.61, 95% confidence interval = 2.25 to 5.78; P<.001, log-rank test). Conclusions: Gene expression grade index appeared to reclassify patients with histologic grade 2 tumors into two groups with high versus low risks of recurrence. This approach may improve the accuracy of tumor grading and thus its prognostic value.

Source

unt

http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE2990 http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE6532

References

Christos Sotiriou and Pratyaksha Wirapati and Sherene Loi and Adrian Harris and Steve Fox and Johanna Smeds and Hans Nordgren and Pierre Farmer and Viviane Praz and Benjamin Haibe-Kains and Christine Desmedt and Denis Larsimont and Fatima Cardoso and Hans Peterse and Dimitry Nuyten and Marc Buyse and Marc J. Van de Vijver and Jonas Bergh and Martine Piccart and Mauro Delorenzi (2006) "Gene Expression Profiling in Breast Cancer: Understanding the Molecular Basis of Histologic Grade To Improve Prognosis", *Journal of the National Cancer Institute*, **98**(4):262-272

Examples

load Biobase package library(Biobase) ## load the dataset data(unt) ## show the first 5 rows and columns of the expression data exprs(unt)[1:5,1:5] ## show the first 6 rows of the phenotype data head(pData(unt)) ## show first 20 feature names featureNames(unt)[1:20] ## show the experiment data summary experimentData(unt) ## show the used platform annotation(unt) ## show the abstract for this dataset abstract(unt)

Index

* datasets unt, 2

unt, <mark>2</mark>